

# Carbendazim

Carbendazim is absorbed through the roots and leaves with acropetal translocation in the plant. It controls *Fungi imperfecti*, *Ascomycetes* and *Basidiomycetes* in a wide range of crops. Carbendazim is used for foliar as well as seed treatments.

HELM provides Carbendazim as **biocide** for industrial use/conservation.



**Action:**

- systemic
- protective and curative

**FRAC classification:** 1

**CAS No.:** 10605-21-7

**Technical grade:** min. 98 %

- white/grey

**Formulations:** 500 SC, 50 WP

**Packaging<sup>1</sup>:**

- 25 kg (tech.)
- 200, 19, 10, 5, 1 ltr. (form. SC)
- 25, 5, 1 kg (form. WP)

Crops	Diseases	Dose rate <sup>2</sup> g a.i./100 ltr.
Apple	Scabs ( <i>Venturia</i> ssp.)	13 – 20
Cereals	Powdery mildew ( <i>Erysiphe graminis</i> ), Stem rot ( <i>Pseudocercospora herpotrichoides</i> ), <i>Fusarium</i> ssp., <i>Septoria</i> ssp.	110 – 210/ha
Citrus	Green mould ( <i>Penicillium digitatum</i> ), Scab ( <i>Sphaceloma fawcetti</i> )	20 – 40
Grapes	Powdery mildew ( <i>Uncinula necator</i> ), <i>Botrytis</i> ssp.	20 – 40
Rice	Rice blast ( <i>Pyricularia oryzae</i> )	110 – 210/ha
Stone fruit	Rots ( <i>Monilia fructigena</i> , <i>Sclerotinia sclerotiorum</i> )	25
Tomato	Collar rot ( <i>Sclerotinia sclerotiorum</i> ), Grey mould ( <i>Botrytis cinerea</i> , <i>Phoma lycopersici</i> )	25

<sup>1</sup>upon request we provide customized packaging

<sup>2</sup>always follow national label recommendations

