

Fosetyl-AI

Fosetyl-AI is a broad-spectrum fungicide that acts by inhibiting germination of spores or by blocking development of mycelium and sporulation. It is rapidly absorbed by leaves and roots with both acropetal and basipetal translocation. The product is often used to control downy mildews and rots in a range of fruit and vegetable crops.



- Action:**
- systemic
 - protective and curative

FRAC classification: 33

CAS No.: 39148-24-8

Technical grade: min. 95 %

Formulations: 80 WP, 80 WG

- Packaging¹:**
- 25 kg (tech.)
 - 25, 1 kg and 100 g sachets (form.)

Crops	Diseases	Dose rate ² g a.i./ha
Citrus	Gummosis – root & foot rots (<i>Phytophthora citrophthora</i>), Brown spots (<i>Alternaria</i> spp.)	Root dip: 2800 Foliar: 4700 Chemigation: 4500 Trunk spray: 40000 –
Pome fruit	Root and collar rots (<i>Phytophthora</i> spp.), Fire blight (<i>Erwinia amylovora</i>)	80000 Foliar: 40000 – 80000 Root dip: 3000
Stone fruit	Root and collar rots (<i>Phytophthora</i> spp.), Almond pruning-wound canker (<i>Pseudomonas syringae</i>)	2200 – 4500 2800 – 4800
Strawberry	Red root rot (<i>Phytophthora fragariae</i>), Leather rot (<i>Phytophthora cactorum</i>)	Foliar: 2200 – 4500 Dip: 2200
Tomato	Root rots (<i>Phytophthora</i> spp.), Dumping off (<i>Pythium debaryanum</i>)	2200 – 4500
Vegetables	Downy mildews (<i>Peronospora</i> spp., <i>Bremia lactucae</i>), Purple blotch (<i>Alternaria porri</i>)	2000 – 4500

¹upon request we provide customized packaging

²always follow national label recommendations

